



# Construction of a High-Density Genetic Map Based on Large-Scale Marker Development in Mango Using Specific-Locus Amplified Fragment Sequencing (SLAF-seq)

Chun Luo <sup>†</sup>, Bo Shu <sup>†</sup>, Quangsheng Yao, Hongxia Wu, Wentian Xu and Songbiao Wang <sup>\*</sup>

Key Laboratory of Tropical Fruit Biology, Ministry of Agriculture; South Subtropical Crops Research Institute, Chinese Academy of Tropical Agricultural Sciences, Zhanjiang, China

## OPEN ACCESS

### Edited by:

Claudio Bonghi,  
University of Padua, Italy

### Reviewed by:

Shengwu Hu,  
Northwest A&F University, China  
Stefano Tartarini,  
University of Bologna, Italy

### \*Correspondence:

Songbiao Wang  
wsbcjy@163.com

<sup>†</sup>These authors have contributed  
equally to this work.

### Specialty section:

This article was submitted to  
Crop Science and Horticulture,  
a section of the journal  
Frontiers in Plant Science

**Received:** 09 May 2016

**Accepted:** 16 August 2016

**Published:** 30 August 2016

### Citation:

Luo C, Shu B, Yao Q, Wu H, Xu W  
and Wang S (2016) Construction of a  
High-Density Genetic Map Based on  
Large-Scale Marker Development in  
Mango Using Specific-Locus  
Amplified Fragment Sequencing  
(SLAF-seq). *Front. Plant Sci.* 7:1310.  
doi: 10.3389/fpls.2016.01310

Genetic maps are particularly important and valuable tools for quantitative trait locus (QTL) mapping and marker assisted selection (MAS) of plant with desirable traits. In this study, 173 F<sub>1</sub> plants from a cross between *Mangifera indica* L. “Jin-Hwang” and *M. indica* L. “Irwin” and their parent plants were subjected to high-throughput sequencing and specific-locus amplified fragment (SLAF) library construction. After preprocessing, 66.02 Gb of raw data containing 330.64 M reads were obtained. A total of 318,414 SLAFs were detected, of which 156,368 were polymorphic. Finally, 6594 SLAFs were organized into a linkage map consisting of 20 linkage groups (LGs). The total length of the map was 3148.28 cM and the average distance between adjacent markers was 0.48 cM. This map could be considered, to our knowledge, the first high-density genetic map of mango, and might form the basis for fine QTL mapping and MAS of mango.

**Keywords:** *Mangifera indica* L., SLAF-seq, high-density genetic map, SNPs, linkage group

## INTRODUCTION

Mango (*Mangifera indica* L.,  $2n = 40$ ) is a major fruit crop of the tropics and subtropics, and is often called the “king of fruits” (Mukherjee, 1950, 1957; Purseglove, 1972). Mango has been referred to as an allopolyploid (Mukherjee, 1950), but the recent microsatellite marker studies of Duval et al. (2005), Viruel et al. (2005), and Schnell et al. (2005) indicate that *M. indica* is diploid. Although mango has a high level of heterozygosity, it has a relatively small genome size (approximately 439 Mb; Arumuganathan and Earle, 1991). In recent years, the area under mango plantation has expanded unceasingly worldwide, and mango yields have increased year by year. Currently, more than 100 countries and regions worldwide produce mango, and China is one of the major mango-producing countries. The mango industry has become the mainstay of the local agricultural industry in most of the mango production provinces in China.

Mango breeding has gained increasing attention in recent years, and conventional breeding methods (cross breeding, mutation breeding, and seedling selection) are mainly used for creating new varieties of mango. However, these methods are time-consuming and difficult to improve upon because of some inherent characteristics of mango (Iyer and Schnell, 2009). Many important economic traits of mango are controlled by a number of loci. Quantitative trait locus (QTL) analysis and marker assisted selection (MAS) can shorten selection times and accelerate the

breeding process of new varieties of mango. Genetic maps, especially high-density genetic maps, are particularly valuable tools for QTL mapping and MAS. Four genetic maps of mango have been generated to date. Chunwongse et al. (2000) first constructed a genetic map of mango using 197 restriction fragment length polymorphism (RFLP) and 650 amplified fragment length polymorphism (AFLP) markers based on 31 F<sub>1</sub> plants from a cross between Alphonso and Palmer cultivars of mango (*M. indica*). Kashkush et al. (2001) developed the second genetic map of mango using 34 AFLP markers based on 29 F<sub>1</sub> individuals and covering a length of 161.5 cM with an average marker spacing of 4.75 cM. Fang et al. (2003) developed the third genetic map of mango using 81 AFLP markers based on 60 F<sub>1</sub> individuals and covering a length of 354.1 cM with an average marker spacing of 4.37 cM. Chunwongse et al. (2015) constructed a partial genetic linkage map spanning a distance of 529.9 cM and consisting of 9 microsatellite and 67 RFLP markers using the same plant materials as in their previous report (Chunwongse et al., 2000). However, these maps were developed on the basis of either a small mapping population (60 or fewer) or a limited number of markers, resulting in a relatively low-density genetic map for future QTL analysis.

DNA markers such as AFLP, RFLP, random amplification of polymorphic DNA (RAPD), inter simple sequence repeat (ISSR), sequence-related amplified polymorphism (SRAP), and simple sequence repeat (SSR) have been developed in mango. However, the current number of markers is too small to build a high-density genetic map. Specific-locus amplified fragment sequencing (SLAF-seq) technology is an efficient method of *de novo* single nucleotide polymorphism (SNP) discovery and large-scale genotyping, which is based on reduced-representation library (RRL) and high-throughput sequencing (Sun et al., 2013). The efficiency of SLAF-seq was tested using rice and soybean data, and has been used to construct the highest-density genetic map of common carp, without a reference genome sequence. To date, SLAF-seq has been used successfully to construct high-density genetic maps and study the genomes of many crops (Chen et al., 2013; Huang et al., 2013; Zhang et al., 2013, 2015; Li et al., 2014; Qi et al., 2014; Wei et al., 2014; Guo et al., 2015; Jiang et al., 2015; Xu et al., 2015; Zhu et al., 2015).

We conducted a hybridization breeding study of mango 10 years ago and established many segregating populations. Following several years' field investigation, we selected an intraspecific cross of *M. indica* for constructing its genetic linkage map. We employed the recently developed SLAF-seq approach to identify a large number of SNP markers for mango and thereby developed a high-density genetic map of mango. The characteristics and value of this genetic map were analyzed and discussed.

## MATERIALS AND METHODS

### Plant Materials

The F<sub>1</sub> mapping population consisted of 173 individuals from a cross between *M. indica* "Jin-Hwang" (female parent) and *M. indica* "Irwin" (male parent) grown at the South Subtropical Crops Research Institute, Zhanjiang, China. The Jin-Hwang

mango has a large, high-quality, and greenish yellow fruit. An important trait of the Jin-Hwang mango is its resistance to anthracnose, to which it is exposed frequently during its cultivation and postharvest storage, and severely affects the development of the mango industry (Lei et al., 2006). The Irwin mango has a medium-sized, high-quality bright yellow fruit with a crimson red blush. However, it is susceptible to anthracnose (Campbell, 1992; Lei et al., 2006).

Young leaves from the parents and F<sub>1</sub> individuals were collected and genomic DNA was isolated by the cetyltrimethylammonium bromide (CTAB) method (Kashkush et al., 2001). The genomic DNA was visualized by electrophoresis in agarose gel and quantified using a NanoDrop 2000 Spectrophotometer (Thermo scientific, USA).

### SLAF Library Construction and High-Throughput Sequencing

The SLAF-seq strategy (Sun et al., 2013) was used in this study. The genomic DNA of the two parents and F<sub>1</sub> population was digested using *Hpy166II* restriction enzyme [New England Biolabs (NEB), USA]. Subsequently, Klenow Fragment (3' → 5' exo-) (NEB) and dATP were used to add a single nucleotide (A) overhang to the digested fragments at 37°C. T4 DNA ligase was used to ligate the duplex tag-labeled sequencing adapters (PAGE-purified; Life Technologies, USA) to the A-tailed fragments. PCR was carried out using diluted restriction-ligation DNA samples, Q5<sup>®</sup> High-Fidelity DNA Polymerase (NEB), dNTPs, the forward primer (5'-AATGATACGGCGACCACCGA-3'), and the reverse primer (5'-CAAGCAGAAGACGGCATACG-3'). The PCR products were then purified using Agencourt AMPure XP beads (Beckman Coulter, High Wycombe, UK) and pooled. The pooled samples were separated using 2% agarose gel electrophoresis. Fragments ranging from 264 to 464 bp (with indexes and adaptors) in size were excised and purified using a QIAquick gel extraction kit (Qiagen, Hilden, Germany). Gel-purified products were then diluted. Paired-end sequencing (125 bp from both ends) was performed using an Illumina HiSeq 2500 system (Illumina, Inc., San Diego, CA, USA) according to the manufacturer's instructions.

### Sequence Data Grouping and Genotyping

SLAF marker identification and genotyping were performed using the procedures described by Sun et al. (2013). Low-quality reads (quality score <20 e) were deleted and the raw reads were assigned to 173 individuals samples according to the duplex barcode sequences. After trimming the barcodes and the terminal 5-bp positions from each high-quality read, the clean reads were clustered together according to their sequence identities. Sequences mapping to the same locus with over 90% identity were defined as one SLAF locus (Zhang et al., 2015). SNP loci between the two parents were detected, and the SLAFs with >3 SNPs were removed. Alleles of each SLAF were defined according to parental reads with sequence depth >10-fold and offspring reads with sequence depth >2-fold. As *M. indica* is a diploid species, one SLAF locus can contain maximum 4 genotypes; therefore, the SLAF loci with >4 alleles were eliminated. Only SLAFs with 2, 3, or 4 alleles were found to be polymorphic

and considered potential markers. All polymorphism markers were grouped into eight segregating patterns. As the map was constructed using the  $F_1$  population from two heterozygous parents, the markers from the segregation pattern of  $aa \times bb$  were filtered out.

In order to ensure the genotyping quality, genotype scoring was conducted using a Bayesian approach as described by Sun et al. (2013). Subsequently, three steps were taken to screen the high-quality markers. First, the markers whose average sequence depths were  $<10$ -fold in parents and  $<6$ -fold in progeny were filtered. Second, the markers with  $>10\%$  missing data were removed. Third, the markers with significant segregation distortion ( $P < 0.05$ ) based on the chi-square test were initially excluded from the genetic map construction and then added later as accessory markers.

## Genetic Map Construction

Marker loci were partitioned primarily into linkage groups (LGs) by modified logarithm of odds (MLOD) scores  $>5$ . For efficient map construction, the HighMap strategy was used to arrange the SLAF markers in a specific order and correct genotyping errors within LGs (Liu et al., 2014). The genetic map was constructed according to the maximum likelihood method (van Ooijen, 2011) and the genotyping errors were corrected by the SMOOTH algorithm (van Os et al., 2005). The missing genotypes were imputed by using a k-nearest neighbor algorithm (Huang et al., 2011). The Kosambi mapping function was applied to estimate the genetic map distances in centimorgan (cM; Kosambi, 1943).

## RESULTS

### Analysis of SLAF Sequencing Data and Genotyping

After preprocessing, 66.02 Gb of raw data containing 330.64 M reads were generated. The raw data of SLAF-seq were submitted to the Sequence Read Archive (SRA) database (accession number: SRX1741570) of the National Center of Biotechnology Information (NCBI). On an average, Q30 (quality scores of at least 30, indicating a 1% chance of error) was 85.60% and the GC content was 36.55%. The numbers of reads for female and male parents were 9,403,617 and 10,407,769, respectively. The read numbers for each  $F_1$  individual ranged from 1,178,049 to 2,363,336 with an average of 1,796,718 (Table S1). After read clustering, a total of 318,414 SLAFs were detected, and their average sequencing depth was found to be 25.35-fold and 6.54-fold for parents and each progeny, respectively (Figure 1).

Among the 318,414 SLAFs, 156,368 were polymorphic with a polymorphism rate of 49.11%, while the remaining 162,046 were non-polymorphic or repetitive. After filtering the SLAF markers lacking the parent information, 125,815 polymorphic markers were successfully genotyped and grouped into eight segregation patterns ( $ab \times cd$ ,  $ef \times eg$ ,  $hk \times hk$ ,  $lm \times ll$ ,  $nn \times np$ ,  $aa \times bb$ ,  $ab \times cc$ , and  $cc \times ab$ ; Figure 2). Since the two parents were heterozygous, the markers from the segregation pattern of  $aa \times bb$  were filtered out.

## Basic Characteristics of the Genetic Map

After a series of screenings, 7394 SLAF markers were found to be effective and used for the final linkage analysis. The average integrity of the mapped markers was 99.90%, indicating a relatively high quality of the genetic map. After linkage analysis, among the 7394 markers mapped onto the 20 LGs, 4866 markers were used for the female map, 2585 for the male map, and 6594 for the integrated map (Figure 3, Tables S2–S4). The total genetic lengths of the female, male, and integrated maps were 3144.23, 2747.89, and 3148.28 cM, respectively. The average distances between the adjacent markers in the female, male, and integrated maps were 0.65, 1.07, and 0.48 cM, respectively.

All LGs are shown in Table 1. For the integrated map, the longest LG was LG19 (206.90 cM), which contained 520 SLAF markers, while the shortest LG (LG6, 119.44 cM) harbored 255 SLAF markers. LG16 had the maximum markers (591), whereas LG8 possessed the minimum markers (219). On an average, each LG contained 330 SLAF markers. The genetic length of the LGs ranged from 119.44 cM (LG6) to 206.90 cM (LG19), with an average distance between the adjacent markers ranging from 0.35 cM (LG16) to 0.8 cM (LG8). The “Gap  $\leq 5$ ” value, which reflected the degree of linkage among markers, ranged from 95.15 to 99.49%, with the largest gap of 18.734 cM located in LG9. A total of 13,844 SNP loci were identified among the 6594 mapped SLAF markers (Table 2). LG16 had the maximum SNP loci (1139), whereas LG8 contained the minimum SNPs (443).


## Quality Evaluation of the Genetic Map

Quality of the mango genetic map was evaluated by constructing the haplotype and heat maps. The haplotype maps, which reflect the double exchange of the population, were developed for the parental controls and 173 offsprings using 6594 SLAF markers (Supplementary Material Presentation 1). Most of the recombination blocks were distinctly defined. The missing data for each LG ranged from 0.04% (LG1) to 0.22% (LG8). Most of the LGs were uniformly distributed, suggesting that the genetic maps were of high quality.


The heat maps showed the relationships of recombination between markers from each LG. Pair-wise comparisons between markers were used to assign recombination scores to 6594 markers, after which the heat maps were constructed (Supplementary Material Presentation 2). The resulting maps showed that the order of SLAF markers in most of the LGs was correct.

## Segregation Distortion Markers on the Map

Of the 6594 markers, only 174 (2.64%) exhibited significant segregation distortion ( $P < 0.05$ ) on the genetic map on the basis of a chi-square test (Table 1). The segregation distortion markers were distributed on most of the LGs with the exceptions of LG10, LG17, LG19, and LG20. The frequencies of distorted markers on LG3 (16.09%) and LG5 (12.64%) were higher than those of the other LGs. No significant correlation was observed between the distribution of the distorted and mapped markers. For example, LG16, which possessed the maximum markers (591 SLAF markers) and covered 205.69 cM, included only one distorted marker. Comparatively, LG8, which possessed the



**FIGURE 1 | Number and coverage of valid SLAF markers for each  $F_1$  individual.** The x-axis indicates the plant accession including female parent (F), male parent (M), and each of the  $F_1$  individuals; the y-axis indicates the number of SLAF markers (A) and the coverage (B).




**FIGURE 2 | Number of polymorphic SLAF markers for eight segregation patterns.** The x-axis indicates eight segregation patterns of polymorphic SLAF markers; the y-axis indicates the number of markers.

minimum markers (219 SLAF markers), included 15 distorted markers.

## DISCUSSION

The development of abundant and reliable molecular markers was very important for the construction of a genetic map. In the present study, SNP markers were used for map construction. SLAF-seq technology is an efficient method of *de novo* SNP detection and large-scale genotyping, which is based on RRL and high-throughput sequencing (Sun et al., 2013). The markers developed using SLAF-seq technology were of relatively higher density, better consistency and effectiveness, and lower cost than those developed using traditional methods. Since the development of SLAF-seq technology, it has been applied in many plant studies and has produced remarkable results. Zhang et al. (2013) employed SLAF-seq technology to detect 71,793 high-quality SLAFs, of which 3673 were polymorphic; they





**FIGURE 3 | Distribution of SLAF markers on the 20 linkage groups of mango.** A black bar indicates a SLAF marker. A red bar indicates a segregation distortion marker. The x-axis represents the linkage group number and the y-axis indicates the genetic distance (cM) in each linkage group.

used 1233 of the polymorphic markers to construct the first high-density genetic map for sesame. Qi et al. (2014) applied SLAF-seq to develop 12,577 polymorphic SLAFs, and used 5308 of them to construct a soybean genetic map. Similar studies have been conducted on many other crops, such as wax gourd (22,151 polymorphic SLAFs; Jiang et al., 2015), tea plant (25,014 polymorphic SLAFs; Ma et al., 2015), walnut (49,174 polymorphic SLAFs; Zhu et al., 2015), mei (93,031 polymorphic SLAFs; Zhang et al., 2015), grapevine (42,279 polymorphic SLAFs; Guo et al., 2015), red sage (62,834 polymorphic SLAFs; Liu et al., 2016), and cucumber (15,946 polymorphic SLAFs; Zhu et al., 2016). In this study, 66.02 Gb of raw data containing 330.64 M reads were generated through SLAF-seq. There were a total of 318,414 SLAF markers, of which 156,368 were polymorphic. After a series of screenings, 7394 SLAF markers were found to be effective and were used for linkage analysis; however, only 6594 markers containing 13,844 SNP loci were mapped successfully onto the high-density genetic map. The average sequencing depth of the markers that failed to be included in the map was 535,794, and the average integrity of all samples was 95%, which suggested that this failure was not due to a sequencing problem. The polymorphic markers of mango developed by SLAF-seq technology were more numerous than those of the previously mentioned crops, further demonstrating the potential of SLAF-seq as a low-cost technique to effectively develop numerous and reliable molecular markers for fruit trees.

Genetic maps are the basis of QTL analyses of the agronomic traits, which are important for the improvement of breeding programs. However, the construction of genetic maps is difficult, and there is a shortage of genetic maps for fruit trees, especially the tropical ones. There were a few reports for papaya (Sondur et al., 1996), avocado (Sharon et al., 1997), mango

(Chunwongse et al., 2000, 2015; Kashkush et al., 2001; Fang et al., 2003), and guava (Padmakar et al., 2015) with respect to genetic maps. Although four genetic maps have been previously reported for mango, there were several deficiencies in them. Firstly, the maximum number of  $F_1$  individuals used for these studies was 60. If the size of the population used for map construction is too small, it cannot reflect small recombination rates, and is not beneficial for increasing the efficiency of QTL detection. Secondly, the dominant molecular markers, such as AFLP, were mainly employed, whereas the co-dominant markers, such as SSR, were rarely used. Thirdly, the number of mapped markers was relatively small, which resulted in relatively large average distances between the adjacent markers. The progress of the genetic map construction was very slow, and to the best of our knowledge, no high-density map has been reported to date. The total length of the new genetic map, which was constructed using 6594 SLAF markers in this study, was 3148.28 cM spanning 20 LGs, and the average distance between adjacent markers was 0.48 cM. This map could be considered the first high-density genetic map of mango, and its evaluation using haplotype and heat maps suggested that it was of high quality (**Supplementary Material Presentations 1, 2**).

Some of the seedling trees of the  $F_1$  progenies of the two cultivars of mango began to set fruit two years ago, and the phenotypic data of the two parents and  $F_1$  progenies were collected. The characteristic of disease resistance was segregated. Since the Jin-Hwang mango is resistant to anthracnose and Irwin mango is susceptible to anthracnose, the cross between the two cultivars might be used to breed a new variety with high anthracnose resistance and enable the study of the inheritance of resistance. The high-density genetic map of mango constructed here will be applied to map the QTL/QTLs of this trait.

TABLE 1 | Basic characteristics of the 20 linkage groups of mango.

Linkage group	Total number of markers			Total distance (cM)			Average distance (cM)			Gap < 5 (cM)			Max gap (cM)		Number of markers with segregation distortion
	Female map	Male map	Integrated map	Female map	Male map	Integrated map	Female map	Male map	Integrated map	Female map	Male map	Integrated map	Female map	Male map	
LG1	230	145	362	161.96	113.72	168.04	0.71	0.79	0.47	96.51%	92.36%	98.06%	22.148	22.188	17
LG2	170	111	272	180.35	165.48	173.21	1.07	1.50	0.64	94.67%	93.64%	97.05%	21.295	23.033	8
LG3	245	67	299	198.74	104.87	170.13	0.81	1.59	0.57	96.31%	90.91%	98.32%	13.030	14.758	28
LG4	390	162	526	188.39	137.92	184.04	0.48	0.86	0.35	99.23%	95.65%	99.43%	13.548	31.201	1
LG5	209	80	262	144.75	88.97	150.23	0.70	1.13	0.58	95.67%	93.67%	97.70%	10.94	17.055	22
LG6	207	76	255	146.29	108.31	119.44	0.71	1.44	0.47	98.06%	92.00%	96.85%	16.249	38.316	3
LG7	195	49	224	125.31	116.33	121.40	0.65	2.42	0.54	96.39%	91.67%	97.76%	12.290	39.192	9
LG8	191	74	219	167.78	105.03	175.09	0.88	1.44	0.80	97.37%	91.78%	96.79%	28.000	45.865	15
LG9	202	171	368	182.83	258.06	198.19	0.91	1.52	0.54	95.02%	93.53%	97.28%	20.624	45.959	3
LG10	192	130	287	119.61	122.7	122.33	0.63	0.95	0.43	96.86%	96.12%	99.30%	9.61	34.369	0
LG11	392	235	480	178.46	189.62	184.04	0.46	0.81	0.38	98.21%	96.58%	99.37%	9.523	19.466	12
LG12	160	108	231	136.15	106.6	126.40	0.86	1.00	0.55	93.71%	93.46%	98.70%	12.401	21.295	9
LG13	264	160	371	160.1	141.03	150.57	0.61	0.89	0.41	97.72%	93.08%	98.92%	16.249	21.295	17
LG14	219	86	272	116.84	109.00	123.21	0.54	1.28	0.45	96.33%	96.47%	98.89%	6.143	46.874	14
LG15	183	98	262	140.72	85.10	141.19	0.77	0.88	0.54	95.05%	92.78%	96.17%	9.926	16.249	4
LG16	477	223	591	206.09	169.13	205.69	0.43	0.76	0.35	98.74%	96.40%	99.49%	11.665	31.035	1
LG17	174	80	228	132.75	116.64	133.93	0.77	1.48	0.59	95.38%	86.08%	95.15%	45.959	24.032	0
LG18	158	160	266	140.96	192.00	170.57	0.90	1.21	0.64	93.63%	96.23%	98.11%	21.026	33.952	11
LG19	359	294	520	193.2	258.51	206.90	0.54	0.88	0.40	98.04%	95.22%	98.27%	38.932	27.908	0
LG20	249	76	299	122.95	58.87	123.68	0.50	0.78	0.42	97.18%	88.00%	97.99%	10.227	6.705	0
Total	4866	2585	6594	3144.23	2747.89	3148.28	0.65	1.07	0.48	96.50%	93.28%	97.98%	45.959	46.874	174

\*Gap < 5" indicates the percentages of gaps in which the distance between the adjacent markers was less than 5cM.

**TABLE 2 | Distribution of SNP loci on the 20 linkage groups of mango.**

Linkage group	SNP number	Transition/Transversion number
LG1	750	487/263
LG2	575	371/204
LG3	691	439/252
LG4	1112	692/421
LG5	592	404/189
LG6	561	389/172
LG7	512	340/172
LG8	443	278/165
LG9	775	497/279
LG10	596	421/176
LG11	971	578/393
LG12	474	312/162
LG13	788	523/265
LG14	562	355/207
LG15	506	319/188
LG16	1139	726/414
LG17	515	351/164
LG18	549	329/221
LG19	1105	705/400
LG20	628	397/231

Additionally, our sequencing results provided a mass of SLAF markers for mango. Because they were explored at the whole-genome level, the sequence and location information of these markers can be used as a reference in the future genome assemblies of mango. Furthermore, a total of 13,844 SNPs, which are co-dominant markers, were found among the SLAF markers, and they can be used for comparative genomic studies and future MAS in mango breeding program. Like other molecular markers, the SLAFs and SNPs developed here also can be applied to other mango cultivars and progenies for identification of the germplasm or hybrid and analysis of the genetic diversity among cultivars.

## REFERENCES

- Arumuganathan, K., and Earle, E. D. (1991). Nuclear DNA content of some important plant species. *Plant Mol. Biol. Rep.* 9, 208–218. doi: 10.1007/BF02672069
- Campbell, R. J. (1992). *A Guide to Mangoes in Florida*. Miami, FL: Fairchild Tropical Garden.
- Chen, S., Huang, Z., Dai, Y., Qin, S., Gao, Y., Zhang, L., et al. (2013). The development of 7E chromosome-specific molecular markers for *Thinopyrum elongatum* based on SLAF-seq technology. *PLoS ONE* 8:e65122. doi: 10.1371/journal.pone.0065122
- Chunwongse, C., Phumichai, C., Tongyoo, P., Juejun, N., and Chunwongse, J. (2015). Development of di-nucleotide microsatellite markers and construction of genetic linkage map in mango (*Mangifera indica* L.). *Songklanakarin J. Sci. Technol.* 37, 119–127.
- Chunwongse, J., Phumichai, C., Barbrasert, C., Sukonsawan, S., Boonreungrawd, R., and Babprasert, C. (2000). Molecular mapping of mango cultivars 'Alphonso' and 'Palmer'. *Acta Hort.* 509, 193–206. doi: 10.17660/ActaHortic.2000.509.19
- Duval, M. F., Bunel, J., Sitbon, C., and Risterucci, A. M. (2005). Development of microsatellite markers for mango (*Mangifera*

## AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: CL, BS, and QY. Performed the experiments and analyzed the data: CL, BS, and HW. Wrote the manuscript: CL, BS. Read and approved the final manuscript: WX, SW.

## ACKNOWLEDGMENTS

This research was supported by the Natural Science Foundation of China (No. 31501721), the National Science Foundation of Guangdong Province (No. 2014A030310341), the National Science Foundation of Hainan Province (No. 20153044), and the Special Scientific Research Fund of Agricultural Public Welfare Profession of China (201203092-3).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2016.01310>

**Supplementary Material Presentation 1 | Haplotype map of the integrated maps.** Each row represents a marker. Markers are ranked in accordance with the map order. Each of the two columns represents an individual; blank columns are used between two individuals. The first and second columns represent the paternal and maternal chromosomes, respectively. The green and blue areas in the columns represent the first and second alleles from the parents, respectively. The white column represents the source of alleles that cannot be judged. The gray areas represent the deleted alleles.

**Supplementary Material Presentation 2 | Heat map of the integrated maps.** Markers of each row and column are ranked according to the map order; each small square represents the rate of recombination (*r*) between the two markers.

**Table S1 | Summary of the sequencing data.**

**Table S2 | SLAF markers and SNP loci on the map.** The SNPs within the SLAF marker allele sequences were marked in lowercase letters. The SLAF markers separated by linkage groups are shown in different sheets and the parent from which the alleles were derived are shown.

**Table S3 | Details of marker genotypes for F1 individuals.**

**Table S4 | SLAF markers and their locations in each LG on the map.**

*indica*, L.). *Mol. Ecol. Notes* 5, 824–826. doi: 10.1111/j.1471-8286.2005.01076.x

Fang, J. G., Liu, D. J., and Ma, Z. Q. (2003). Constructing mango (*Mangifera indica* L.) genetic map using markers for double heterozygous loci. *Mol. Plant Breed.* 1, 313–319. (in Chinese). doi: 10.3969/j.issn.1672-416X.2003.03.004

Guo, Y., Shi, G., Liu, Z., Zhao, Y., Yang, X., Zhu, J., et al. (2015). Using specific length amplified fragment sequencing to construct the high-density genetic map for *Vitis* (*Vitis vinifera* L. × *Vitis amurensis* Rupr.). *Front. Plant Sci.* 6:393. doi: 10.3389/fpls.2015.00393

Huang, S., Ding, J., Deng, D., Tang, W., Sun, H., Liu, D., et al. (2013). Draft genome of the kiwifruit *Actinidia chinensis*. *Nat. Commun.* 4:2640. doi: 10.1038/ncomms3640

Huang, X., Zhao, Y., and Wei, X., Li, C., Wang, A., Zhao, Q., et al. (2011). Genome-wide association study of flowering time and grain yield traits in a worldwide collection of rice germplasm. *Nat. Genet.* 44, 32–39. doi: 10.1038/ng.1018

Iyer, C. P. A., and Schnell, R. J. (2009). "Breeding and genetics," in *The Mango: Botany, Production and Uses*, ed R. E. Litz (London: British Library), 67–96.

Jiang, B., Liu, W., Xie, D., Peng, Q., He, X., Lin, Y., et al. (2015). High-density genetic map construction and gene mapping of pericarp color in wax gourd using specific-locus amplified fragment (SLAF) sequencing. *BMC Genomics* 16:1035. doi: 10.1186/s12864-015-2220-y

- Kashkush, K., Fang, J., Eli, T., Hillel, J., and Lavi, U. (2001). Cultivar identification and genetic map of mango (*Mangifera indica*). *Euphytica* 1221, 129–136. doi: 10.1023/A:1012646331258
- Kosambi, D. D. (1943). The estimation of map distances from recombination values. *Ann. Hum. Genet.* 12, 172–175. doi: 10.1111/j.1469-1809.1943.tb02321.x
- Lei, X., Zhao, Y., Yao, Q., He, Y., Sun, G., Ma, W., et al. (2006). Identification and analysis of the resistance of different mango cultivars to anthracnose (*Colletotrichum gloeosporioides*). *J. Fruit Sci.* 23, 838–842. (in Chinese). doi: 10.3969/j.issn.1009-9980.2006.06.012
- Li, B., Tian, L., Zhang, J., Huang, L., Han, F., Yan, S., et al. (2014). Construction of a high-density genetic map based on large-scale markers developed by specific length amplified fragment sequencing (SLAF-seq) and its application to QTL analysis for isoflavone content in *Glycine max*. *BMC Genomics* 15:1086. doi: 10.1186/1471-2164-15-1086
- Liu, D., Ma, C., Hong, W., Huang, L., Liu, M., Liu, H., et al. (2014). Construction and analysis of high-density linkage map using high-throughput sequencing data. *PLoS ONE* 9:e98855. doi: 10.1371/journal.pone.0098855
- Liu, T., Guo, L., Pan, Y., Zhao, Q., Wang, J., and Song, Z. (2016). Construction of the first high-density genetic linkage map of *Salvia miltiorrhiza* using specific length amplified fragment (SLAF) sequencing. *Sci. Rep.* 6:24070. doi: 10.1038/srep24070
- Ma, J., Huang, L., Ma, C., Jin, J., Li, C., Wang, R., et al. (2015). Large-scale SNP discovery and genotyping for constructing a high-density genetic map of tea plant using specific-locus amplified fragment sequencing (SLAF-seq). *PLoS ONE* 10:e0128798. doi: 10.1371/journal.pone.0128798
- Mukherjee, S. K. (1950). Mango: its allopolyploid nature. *Nature* 166, 196–197.
- Mukherjee, S. K. (1957). Cytology of some Malayan species of *Mangifera*. *Cytologia* 22, 239–241.
- Padmakar, B., Kanupriya, C., Latha, P. M., Prashant, K. S., Dinesh, M. R., and Sailaja, D. (2015). Development of SRAP and SSR marker-based genetic linkage maps of guava (*Psidium guajava* L.). *Sci. Hortic.* 192, 158–165. doi: 10.1016/j.scienta.2015.04.031
- Purseglove, J. W. (1972). Mangoes west of India. *Acta Hortic.* 24, 107–174. doi: 10.17660/actahortic.1972.24.30
- Qi, Z., Huang, L., Zhu, R., Xin, D., Liu, C., Han, X., et al. (2014). A high-density genetic map for soybean based on specific length amplified fragment sequencing. *PLoS ONE* 9:e104871. doi: 10.1371/journal.pone.0104871
- Schnell, R. J., Olano, C. T., Quintanilla, W. E., and Meerow, A. W. (2005). Isolation and characterization of 15 microsatellite loci from mango (*Mangifera indica* L.) and cross-species amplification in closely related taxa. *Mol. Ecol. Notes* 5, 625–627. doi: 10.1111/j.1471-8286.2005.01018.x
- Sharon, D., Cregan, P. B., Mhameed, S., Kusharska, M., Hillel, M., and Lahav, M. (1997). An integrated genetic linkage map of avocado. *Theor. Appl. Genet.* 95, 911–921.
- Sondur, S. N., Manshardt, R. M., and Stiles, J. I. (1996). A genetic linkage map of papaya based on randomly amplified polymorphic DNA markers. *Theor. Appl. Genet.* 93, 547–553.
- Sun, X., Liu, D., Zhang, X., Li, W., Liu, H., Hong, W., et al. (2013). SLAF-seq: an efficient method of large-scale *de novo* SNP discovery and genotyping using high-throughput sequencing. *PLoS ONE* 8:e58700. doi: 10.1371/journal.pone.0058700
- van Ooijen, J. (2011). Multipoint maximum likelihood mapping in a full-sib family of an outbreeding species. *Genet. Res.* 93, 343–349. doi: 10.1017/S0016672311000279
- van Os, H., Stam, P., Visser, R. G. F., and van Eck, H. J. (2005). SMOOTH: a statistical method for successful removal of genotyping errors from high-density genetic linkage data. *Theor. Appl. Genet.* 112, 187–194. doi: 10.1007/s00122-005-0124-y
- Viruel, M. A., Escribano, P., Barbieri, M., Ferri, M., and Hormaza, J. I. (2005). Fingerprinting, embryo type and geographic differentiation in mango (*Mangifera indica*, L. Anacardiaceae) with microsatellites. *Mol. Breeding* 15, 383–393. doi: 10.1007/s11032-004-7982-x
- Wei, Q., Wang, Y., Qin, X., Zhang, Y., Zhang, Z., Wang, J., et al. (2014). An SNP-based saturated genetic map and QTL analysis of fruit-related traits in cucumber using specific-length amplified fragment (SLAF) sequencing. *BMC Genomics* 15:1158. doi: 10.1186/1471-2164-15-1158
- Xu, X., Xu, R., Zhu, B., Yu, T., Qu, W., Lu, L., et al. (2015). A high-density genetic map of cucumber derived from specific length amplified fragment sequencing (SLAF-seq). *Front. Plant Sci.* 5:768. doi: 10.3389/fpls.2014.00768
- Zhang, J., Zhang, Q., Cheng, T., Yang, W., Pan, H., and Zhong, J. (2015). High-density genetic map construction and identification of a locus controlling weeping trait in an ornamental woody plant (*Prunus mume* Sieb. et Zucc.). *DNA Res.* 22, 183–191. doi: 10.1093/dnares/dsv003
- Zhang, Y., Wang, L., Xin, H., Li, D., Ma, C., Ding, X., et al. (2013). Construction of a high-density genetic map for sesame based on large scale marker development by specific length amplified fragment (SLAF) sequencing. *BMC Plant Biol.* 13:141. doi: 10.1186/1471-2229-13-141
- Zhu, Y., Huang, L., Chen, L., Yang, J., Wu, J., Qu, M., et al. (2016). A high-density genetic linkage map for cucumber (*Cucumis sativus* L.): based on specific length amplified fragment (SLAF) sequencing and QTL analysis of fruit traits in cucumber. *Front. Plant Sci.* 7:437. doi: 10.3389/fpls.2016.00437
- Zhu, Y., Yin, Y., Yang, K., Li, J., Sang, Y., Huang, L., et al. (2015). Construction of a high-density genetic map using specific length amplified fragment markers and identification of a quantitative trait locus for anthracnose resistance in walnut (*Juglans regia* L.). *BMC Genomics* 16:614. doi: 10.1186/s12864-015-1822-8

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Luo, Shu, Yao, Wu, Xu and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.